

AD-A034 485

WISCONSIN UNIV MADISON MATHEMATICS RESEARCH CENTER  
COUPLED-OSCILLATOR NATURAL ORBITALS. (U)

F/G 12/1

NOV 76 P D ROBINSON

DAA629-75-C-0024

UNCLASSIFIED

MRC-TSR-1696

NL

[OF]

AD  
A034485



END

DATE  
FILMED

2-77

ADA 034485

MRC Technical Summary Report #1696

COUPLED-OSCILLATOR NATURAL  
ORBITALS

Peter D. Robinson

**Mathematics Research Center  
University of Wisconsin-Madison  
610 Walnut Street  
Madison, Wisconsin 53706**

November 1976

(Received October 19, 1976)



**Approved for public release  
Distribution unlimited**

Sponsored by

U. S. Army Research Office  
P. O. Box 12211  
Research Triangle Park  
North Carolina 27709

University of Wisconsin-Madison  
Graduate School  
Madison, Wisconsin 53706

UNIVERSITY OF WISCONSIN - MADISON  
MATHEMATICS RESEARCH CENTER

COUPLED-OSCILLATOR NATURAL ORBITALS

Peter D. Robinson

Technical Summary Report #1696  
November 1976

ABSTRACT

The natural orbitals are derived for a  
pair of identical coupled harmonic oscillators in  
their ground state.

AMS (MOS) Subject Classifications: 33A65, 45C05, 81.33

Key Words: Natural orbitals, Coupled harmonic oscillators,  
Integral operator, Eigenvectors

Work Unit Number 1 (Applied Analysis)

ACCESSION for	
NTIS	White Section <input checked="" type="checkbox"/>
DDC	Buff Section <input type="checkbox"/>
UNANNOUNCED	<input type="checkbox"/>
JUSTIFICATION.....	
BY.....	
DISTRIBUTION/AVAILABILITY CODES	
SAC. AVAIL. OR SPECIAL	
<i>A</i>	

# COUPLED-OSCILLATOR NATURAL ORBITALS

Peter D. Robinson

Examples of natural orbitals in closed form seem rare enough [1] to justify this note, in which such orbitals are exhibited for coupled oscillators.

Let  $\{\theta_n(\alpha, x)\}$  denote the complete set of orthonormal eigenfunctions of the oscillator Hamiltonian

$$h(\alpha, x) = -\frac{1}{2} \frac{d^2}{dx^2} + \frac{1}{2} \alpha^4 x^2, \quad \alpha > 0, \quad -\infty < x < \infty, \quad (1)$$

corresponding to the eigenvalues  $(n + \frac{1}{2})\alpha^2$ . Specifically

$$\theta_n(\alpha, x) = \left[ \frac{\alpha}{\pi^{\frac{1}{2}} 2^n n!} \right]^{\frac{1}{2}} H_n(\alpha x) \exp(-\frac{1}{2} \alpha^2 x^2), \quad n = 0, 1, 2, \dots, \quad (2)$$

where the Hermite polynomials  $H_n(\alpha x)$  are orthogonalized with respect to a weight factor [2]  $\exp(-\alpha^2 x^2)$ . Consider now the identical coupled oscillators described by the Hamiltonian

$$H = -\frac{1}{2} \frac{d^2}{dx_1^2} - \frac{1}{2} \frac{d^2}{dx_2^2} + \frac{1}{2} a^4 (x_1^2 + x_2^2) - \frac{1}{4} (a^4 - b^4) (x_1 - x_2)^2, \quad (3)$$

$$a > b > 0, \quad -\infty < x_1 < \infty, \quad -\infty < x_2 < \infty.$$

In terms of the coordinates

$$y_1 = \frac{1}{\sqrt{2}} (x_1 + x_2), \quad y_2 = \frac{1}{\sqrt{2}} (x_1 - x_2), \quad (4)$$



the Hamiltonian (3) takes the separable form

$$H = h(a, y_1) + h(b, y_2) . \quad (5)$$

The ground-state wave-function thus has a space part

$$\begin{aligned} \psi(x_1, x_2) &= \theta_0(a, y_1) \theta_0(b, y_2) = \left( \frac{ab}{\pi} \right)^{\frac{1}{2}} \exp\left(-\frac{1}{2} a^2 y_1^2 - \frac{1}{2} b^2 y_2^2\right) \\ &= \left( \frac{ab}{\pi} \right)^{\frac{1}{2}} \exp\left\{-\frac{1}{4} (a^2 + b^2)(x_1^2 + x_2^2) - \frac{1}{2} (a^2 - b^2)x_1 x_2\right\} , \end{aligned} \quad (6)$$

which is the eigenfunction of the operator (5) corresponding to the eigenvalue  $\frac{1}{2} (a^2 + b^2)$ .

This function  $\psi(x_1, x_2)$  is symmetric in  $x_1$  and  $x_2$ , and is normalized so that

$$\int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \psi^2 dx_1 dx_2 = 1 . \quad (7)$$

Accordingly,  $\psi(x_1, x_2)$  is the Hilbert-Schmidt-type kernel of a compact, self-adjoint integral operator on the real Hilbert space  $L_2(-\infty, \infty)$ . Standard integral equation theory [3] tells us that this operator has a complete orthonormal set of eigenvectors  $\{\phi_n\}$  defined by

$$\int_{-\infty}^{\infty} \psi(x_1, x_2) \phi_n(x_2) dx_2 = \lambda_n \phi_n(x_1), \quad n = 0, 1, 2, \dots , \quad (8)$$

with the eigenvalues  $\lambda_n$  accumulating at zero. Further, the expansion

$$\frac{1}{2} \gamma(x_1, x_2) = \int_{-\infty}^{\infty} \psi(x_1, s) \psi(x_2, s) ds = \sum_{n=0}^{\infty} \lambda_n^2 \phi_n(x_1) \phi_n(x_2) \quad (9)$$

is convergent for all  $x_1$  and  $x_2$ . The functions  $\phi_n(x)$  are precisely the natural orbitals for  $\psi(x_1, x_2)$ , and (9) gives the expansion for the first-order density matrix  $\gamma(x_1, x_2)$ . The "occupation numbers"  $\lambda_n^2$  must sum to unity, from (7) and (9). The expansion for  $\psi(x_1, x_2)$  itself, namely

$$\psi(x_1, x_2) \sim \sum_{n=0}^{\infty} \lambda_n \phi_n(x_1) \phi_n(x_2), \quad (10)$$

must also be convergent, at least in the  $L_2$  norm sense.

In the case of the function given in (6), the expansion (10) can be found directly, and is convergent for all  $x_1$  and  $x_2$ . Mehler's formula [2] states that

$$\exp\left\{-\frac{z^2}{1-z^2}(t_1^2 + t_2^2) + \frac{2z}{1-z^2}t_1t_2\right\} = (1-z^2)^{\frac{1}{2}} \sum_{n=0}^{\infty} \frac{z^n}{2^n n!} H_n(t_1) H_n(t_2),$$

$$-1 < z < 1, \quad -\infty < t_1 < \infty, \quad -\infty < t_2 < \infty. \quad (11)$$

Setting  $t_1 = \alpha x_1$ ,  $t_2 = \alpha x_2$  and multiplying through by  $\exp\{-\frac{1}{2}\alpha^2(x_1^2 + x_2^2)\}$ , we obtain from (2):

$$\frac{\alpha}{\pi^{\frac{1}{2}}} \exp\left\{-\left(\frac{1+z^2}{1-z^2}\right) \frac{\alpha^2}{2}(x_1^2 + x_2^2) + 2 \frac{\alpha^2 z}{1-z^2} x_1 x_2\right\} = (1-z^2)^{\frac{1}{2}} \sum_{n=0}^{\infty} z^n \theta_n(\alpha, x_1) \theta_n(\alpha, x_2). \quad (12)$$

With the choices

$$\alpha = (ab)^{\frac{1}{2}}, \quad z = -\frac{a-b}{a+b} \quad (13)$$

the left-hand side of (12) becomes  $\psi(x_1, x_2)$  as given in (6). Thus

$$\psi(x_1, x_2) = \sum_{n=0}^{\infty} \left\{ \frac{2\sqrt{ab}}{a+b} (-1)^n \left( \frac{a-b}{a+b} \right)^n \right\} \theta_n(\sqrt{ab}, x_1) \theta_n(\sqrt{ab}, x_2), \quad (14)$$

so that for the ground-state of the coupled oscillators we have the natural orbitals

$$\phi_n(x) = \theta_n(\sqrt{ab}, x) = \left[ \frac{\sqrt{ab}}{\pi^{\frac{1}{2}} 2^n n!} \right]^{\frac{1}{2}} H_n(\sqrt{ab} x) \exp(-\frac{1}{2} ab x^2), \quad (15)$$

and corresponding eigenvalues

$$\lambda_n = \frac{2\sqrt{ab}}{a+b} (-1)^n \left( \frac{a-b}{a+b} \right)^n. \quad (16)$$

The orbitals (15), and the occupation numbers  $\lambda_n^2$ , can also be obtained by working from the expression for  $\frac{1}{2} \gamma(x_1, x_2)$ , which is

$$\frac{1}{2} \gamma(x_1, x_2) = ab \left[ \frac{\pi}{2} (a^2 + b^2) \right]^{-\frac{1}{2}} \exp \left\{ -\frac{(a^4 + b^4 + 6a^2 b^2)}{8(a^2 + b^2)} (x_1^2 + x_2^2) + \frac{(a^2 - b^2)^2}{4(a^2 + b^2)} x_1 x_2 \right\}. \quad (17)$$

Extension to three-dimensional oscillators, for which Davidson [1] has discussed the first-order density matrix, is straightforward.

#### REFERENCES

1. E. R. Davidson, *Reduced Density Matrices in Quantum Chemistry*, (Academic Press) 1976.
2. A. Erdélyi (Ed.), *Higher Transcendental Functions*, II, Ch. 10, (McGraw-Hill) 1953.
3. R. Courant and D. Hilbert, *Methods of Mathematical Physics*, I, (Interscience) 1953.



UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM	
1. REPORT NUMBER (14) MRC-TSR-1696 ✓	2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER (9) Technical	
4. TITLE (and Subtitle) (6) COUPLED-OSCILLATOR NATURAL ORBITALS. ✓		5. TYPE OF REPORT & PERIOD COVERED Summary Report, no specific reporting period	
7. AUTHOR(s) (10) Peter D. Robinson		6. PERFORMING ORG. REPORT NUMBER	
9. PERFORMING ORGANIZATION NAME AND ADDRESS Mathematics Research Center, University of Wisconsin 610 Walnut Street Madison, Wisconsin 53706 ✓		8. CONTRACT OR GRANT NUMBER(s) (15) DAAG29-75-C-0024 ✓	
11. CONTROLLING OFFICE NAME AND ADDRESS See Item 18 below. (12) 7p.		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS	
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		12. REPORT DATE (11) November 1976	
		13. NUMBER OF PAGES 4	
		15. SECURITY CLASS. (of this report) UNCLASSIFIED	
		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE	
16. DISTRIBUTION STATEMENT (of this Report)  Approved for public release; distribution unlimited.			
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)			
18. SUPPLEMENTARY NOTES U. S. Army Research Office      University of Wisconsin-Madison P. O. Box 12211      Graduate School Research Triangle Park      Madison, Wisconsin 53706 North Carolina 27709			
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Natural orbitals Coupled harmonic oscillators Integral operator Eigenvectors			
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) The natural orbitals are derived for a pair of identical coupled harmonic oscillators in their ground state.			

DD FORM 1 JAN 73 1473

EDITION OF 1 NOV 65 IS OBSOLETE

UNCLASSIFIED 22/ 200  
SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)